



MNP Report 500080007/2007

The Netherlands Informative Inventory Report 2007

B.A. Jimmink, P.G. Ruysenaars, L.J. Brandes, P.W.H.G. Coenen¹,
A. Hoen, J. Jonker¹, C.J. Peek, M.W. van Schijndel, D. Wever

Contact:

Benno Jimmink

Netherlands Environmental Assessment Agency

European Air Quality and Sustainability

benno.jimmink@mnp.nl

¹⁾ TNO Built Environment and Geosciences

This investigation has been performed by order and for the account of Netherlands Ministry of Housing, Spatial Planning, and the Environment, within the framework of MNP project 500080, project title 'Emission Inventory'.

© MNP 2007

Parts of this publication may be reproduced, on condition of acknowledgement: 'Netherlands Environmental Assessment Agency, Netherlands Informative Inventory Report 2007.'

Rapport in het kort

Informatief Inventarisatie Rapport Nederland 2007

Dit Informatief Inventarisatie Rapport (IIR) geeft een toelichting op de officiële emissiecijfers die Nederland heeft geleverd aan het UNECE-secretariaat in het kader van de verplichtingen onder de Convention on Long-range Transboundary Air Pollution (CLRTAP), en aan de Europese Commissie in het kader van de verplichtingen onder de NEC¹-richtlijn. De emissiecijfers zijn te vinden op de EMEP²-website: <http://www.emep.int> (EMEP data).

Op dit moment stellen landen het IIR nog op vrijwillige basis op. Dat zal bij de revisie van de protocollen onder CLRTAP en de NEC-richtlijn van de EU hoogstwaarschijnlijk veranderen. Het belang van de verplichting van het IIR is dat landen meer inzicht geven in de manier waarop de emissies worden berekend.

De IIR-rapportage 2007 biedt een beter zicht op de toepasbaarheid (vergelijkbaarheid tussen landen, modelberekeningen voor luchtkwaliteit door EMEP) en afrekenbaarheid (transparantie, compleetheid, consistentie tussen jaren, sterkte en zwaktes in methoden, onzekerheden) van emissiecijfers.

¹ National Emissions Ceilings Directive.

² Co-operative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (EMEP).

Trefwoorden:

emissieregistratie, LRTAP, NEC, emissie

Contents

Summary	7
1 Introduction	9
1.1 National inventory background.....	9
1.2 Institutional arrangements for inventory preparation.....	10
1.3 The process of inventory preparation.....	10
1.4 Methods and data sources	12
1.5 Key source analysis.....	13
1.6 Reporting, QA/QC and archiving	15
1.7 Uncertainties	16
1.7.1 Quantitative uncertainty	16
1.8 Explanation on the use of notation keys	18
1.9 Missing sources.....	20
2 Trends in emissions	21
2.1 Trends in national emissions.....	21
2.2 Trends for sulphur dioxide (SO ₂).....	22
2.3 Trends for nitrogen oxides (NO _x)	23
2.4 Trends for ammonia (NH ₃)	24
2.5 Trends for non-methane volatile organic compounds (NMVOC).....	25
2.6 Trends for particulate matter (PM ₁₀).....	26
2.7 Trends for heavy metals (Pb and Cd)	28
2.8 Trends for polycyclic aromatic hydrocarbons (PAH) and dioxins	30
3 Energy, stationary fuel combustion (1A)	33
3.1 Energy industries (1A1).....	34
3.2 Manufacturing industries and construction (1A2)	34
3.3 Other sectors (1A4).....	35
3.4 Other (1A5).....	38
3.5 Mobile combustion (1A3).....	39
3.6 Evaporation, tyre and brake wear, road abrasion (1A3b)	42
3.7 Energy, fugitive emissions from fuels (1B).....	44
4 Industry (2)	45
4.1 Mineral production (2A)	46

4.2	Chemical industry (2B)	46
4.3	Metal production (2 C)	47
4.4	Pulp and paper production (2D1)	48
4.5	Food and drink production (2D2)	48
4.6	Other production (2G)	48
5	Solvents and product use (3)	49
5.1	Paint application (3A)	49
5.2	Degreasing and dry cleaning (3B)	49
5.3	Chemical products, manufacture and processing (3C)	49
5.4	Other, including products containing HMs and POPs (3D)	49
6	Agriculture (4)	51
6.1	Dairy cattle (4B1A)	52
6.2	Non-dairy cattle (4B1b)	52
6.3	Swine (4B8)	52
6.4	Poultry (4B9)	53
6.5	Other agricultural emissions (4G)	53
7	Waste (6)	55
7.1	Waste incineration (6C)	55
7.2	Other waste (6D)	55
8	Other (7)	57
9	Recalculations and other changes	59
9.1	Recalculations compared to 2006 submission	59
9.2	Developments in emission insights and estimates	60
9.3	Improvements	62
	References	65

Summary

This report constituting the Netherlands Informative Inventory Report (IIR) contains information on the inventories in the Netherlands up to 2005 (see EMEP¹ data on <http://www.emep.int>). It includes descriptions of methods and data sources, QA/QC activities carried out and a trend analysis.

This IIR outlines such methods for estimating emissions as the extrapolation of emissions from individual companies to sectors. Estimations are given in more detail for sector and subsector for the key source categories (e.g. emission calculation from road transport in vehicle categories and road types).

The 2007 submission includes emission data from the Netherlands for the years 1990 up to and including 2005. The emission data, with the exception of PM_{2.5} emissions, are extracted from the Dutch Emission Inventory system (PER). These data are calculated from the PM₁₀ data and have not yet been incorporated in the PER.

In the 1990 – 2005 period emissions of all gases presented in this report showed a downward trend. The major overall drivers for this trend are emission reductions in the industrial sectors, cleaner fuels and cleaner cars.

Based on methodological improvements (such as improvement of activity data), the historical data for 1990, 1995, 2000 and 2003-2005 are recalculated annually in the Dutch inventory. Data for other years (1991-1994, 1996-1999, 2001 and 2002) have been based on interpolations.

¹ Co-operative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (EMEP).

1 Introduction

Reporting emission data to the Executive Body of the Convention on Long-range Transboundary Air Pollution (CLRTAP) is required to fulfil obligations in compliance with the implementation of Protocols under the Convention. Parties are required to submit reports on annual national emissions of SO₂, NO_x, NMVOC, CO and NH₃, and various heavy metals and POPs using the Guidelines for Estimating and Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution (UNECE, 2003).

The Netherlands Informative Inventory Report (IIR) 2007 contains information on the Netherlands' inventories for the year 2005, including descriptions of methods, data sources, QA/QC activities carried out and a trend analysis. The inventory covers all anthropogenic emissions to be reported in the Nomenclature for Reporting (NFR), except for individual PAHs (with only total emissions reported), which are to be reported under POPs in Annex III. The publication of an IIR is part of the inventory improvement programme.

1.1 National inventory background

Emissions in the Netherlands are registered in the Pollutant Emission Register (PER), the national database for target group monitoring used by the Dutch government to monitor greenhouse gas emissions in conformance with UNFCCC requirements and the Kyoto Protocol (National System). The database is also employed to monitor pollutants within the framework of National Emission Ceilings (EU) and the Convention on Long-range Transboundary Air Pollution (CLRTAP). PER encompasses the process of data collection, data processing, registration and reporting on emission data for some 170 policy-relevant compounds, and compound groups in air, water and soil.

Emission estimates are based mainly on official statistics of the Netherlands, e.g. energy and agricultural statistics, environmental reports of companies in the industrial sector and emission factors (nationally developed factors and internationally recommended ones).

The Netherlands uses the 'Guidelines for Estimating and Reporting Emission Data' for reporting to the Economic Commission for Europe (UNECE) Convention on Long-range Transboundary Air Pollution (CLRTAP). However, instead of using the EMEP/CORINAIR Emission Inventory Guidebook (EEA, 2005), the Netherlands usually applies country-specific methods, including monitoring data and emission factors.

Note that definitions for emissions accounting differ according to the guideline (EMEP/Corinair, UNFCCC, EU).

1.2 Institutional arrangements for inventory preparation

The Dutch Ministry of Housing, Spatial Planning and the Environment (VROM) has the overall responsibility for the emission inventory and submissions to CLRTAP. A Pollutant Emission Register (PER) system has been in operation in the Netherlands since 1974. Since April 2004 the Ministry of VROM has outsourced the full coordination of the PER to the Emission Registration team (ER) at the Netherlands Environmental Assessment Agency (MNP). This has resulted in a clearer definition and distinction between responsibilities, as well as a concentration of tasks.

The main objective of the emission inventory is to produce an annual set of unequivocal emission data, which is up-to-date, complete, transparent, comparable, consistent and accurate. Emission data are produced in an annual (project) cycle (MNP, 2006) and various external agencies contribute to the PER by performing calculations or submitting activity data (see next section). Besides the Netherlands Environmental Assessment Agency (MNP), the following institutes contribute to the PER:

- Statistics Netherlands (CBS);
- TNO Built Environment and Geosciences (TNO);
- Institute for Inland Water Management and Waste Water Treatment (RIZA);
- several institutes related to Wageningen University and Research Centre (WUR);
- SenterNovem (Waste management division);
- Agricultural Economics Research Institute (LEI);
- Facilitating Organisation for Industry (FO-I), which coordinates annual environmental reporting by companies.

Each of the contributing institutes has its own responsibility and role in the data collection, emission calculations and quality control. These are laid down in general agreements with MNP and in the annual project plan (MNP, 2006). The Informative Inventory Report (IIR) was prepared by MNP.

1.3 The process of inventory preparation

Data collection

For the collection and processing of data, the PER is organised in task forces according to pre-determined methods. The task forces are formed by sector experts of the participating institutes. Methods are compiled on the basis of the best available scientific views. Changes in scientific views lead to changes in methods, and to recalculation of the historical emissions. The following task forces are recognised:

- task force on agriculture and land use;
- task force on energy, industry and waste;
- task force on traffic;
- task force on water, and,
- task force on the consumer and service sector.

Every year, after collection of the emission data, several quality control checks are performed in the task forces during a yearly 'trend analysis' workshop. After approval by participating institutes, emission data are released for publication. Subsequently, emission data is des-aggregated to regional emission data for national use (e.g. 5x5 km grid data for provinces).

Data storage

In cooperation with the contributing research institutes, emission data are collected and stored in a database managed by the Environmental Assessment Agency. A separate database contains the information from the Environmental Annual Reports.

About 250 companies are legally obliged to submit an Environmental Annual Report (EAR). As from 1 January 2004, companies may submit their EARs electronically (e-EAR). Each of these companies has an emission monitoring and registration system in which the specifications are in agreement with the supervisory authority. The provincial authorities validate and verify the reported emissions. The (e-)EARs are then entered by FO-Industry into the EAR database. In addition, a number of companies are required to report information under the BEES/A legislation. Other companies (about 200) provide emission data voluntarily within the framework of environmental covenants.

Point-source emission data in the EAR database is validated by the task forces. The result is a selection of validated point source emissions and activity data, which are then stored in the ER-I database. The ER-I data is combined with supplementary estimates for Small and Medium-sized Enterprises (SMEs). Several methods are applied for calculating these emissions. TNO has derived emission factors for NO_x emissions from small installations, for instance (Soest-Vercammen et al., 2002), while, for other substances, the Implied Emission Factors (IEFs) derived from the EARs are applied to calculate sector emissions.

Emissions from the ER-I database and collectively estimated industrial as well as non-industrial sources are stored in the ER-C database (see Figure 1.1). The ER-C database, consisting of a large number of geographically distributed emission sources (about 1200), contains a complete record of Dutch emissions for a particular year. Each emission source includes information on the Standard Industrial Classification code (SBI-code) and industrial subsector, separate information in process and combustion emissions, and the relevant environmental compartment and location. These emission sources can be selectively aggregated, for example, by NFR category.




Figure 1.1. The Netherlands Pollutant Emission Register

1.4 Methods and data sources

Methods used in the Netherlands are documented in several reports and in meta-data files, available on www.emissieregistratie.nl. However, most of these reports are only available in Dutch. For greenhouse gases, particulate matter and all emissions related to mobile sources, the documentation has been translated in English. This documentation can be found on www.prtr.nl.

In general, two emission models are used in the Netherlands:

- A model for emissions of large point sources (e.g. large industrial, power plants) that are registered individually and supplemented with emission estimates for the remainder of the companies in a sector (based mainly on IEFs from the individually registered companies). This is the so-called ‘bottom up’ method.
- A model for emissions of diffuse sources (e.g. road transport, agriculture) that are calculated from activity data and emission factors from sectoral emission inventory studies in the Netherlands (e.g. SPIN documents produced by the ‘Cooperation project on industrial emissions’).

The following sections sketch these methods, which are discussed in more detail, by sector or subsector, for the top ten key sources mentioned in chapter 3.

1.5 Key source analysis

More than 95% of national total should be covered for all key source categories. The REPDAB generated key source category list is insufficient for this purpose. Table 1.1 shows the key source categories which cover 95% of the national total for a component.

Table 1.1.a. Key source categories for SO_x, NO_x, NH₃ and NMVOC

Component	SO _x		NO _x		NH ₃		NMVOC	
Key source categories (Sorted from high to low from top to bottom)	1 A 1 b	47.28%	1 A 3 b iii	23.48%	4 B 1 a	28.20%	3 D	15.85%
	1 A 1 a	17.05%	1 A 3 b i	12.77%	4 B 8	23.30%	3 A	14.80%
	1 A 2 c	6.47%	1 A 1 a	12.76%	4 B 1 b	16.20%	1 A 3 b i	14.59%
	1 A 2 a	6.44%	1 A 2 f	5.78%	4 B 9	12.50%	2 G	8.18%
	1 A 2 f	5.36%	1 A 3 b ii	5.52%	4 G	9.40%	2 B 5	5.64%
	1 A 2 b	5.32%	1 A 4 b i	5.52%	7	3.80%	1 A 3 b iv	5.48%
	1 A 4 c ii	2.51%	1 A 4 c ii	5.32%	1 A 3 b i	1.80%	1 A 4 b i	4.88%
	1 A 3 d ii	1.53%	1 A 3 d ii	4.51%			1 B 2 b	3.72%
	2 A 7	1.46%	1 A 2 c	4.34%			1 B 2 a iv	3.61%
	1 A 4 c iii	1.12%	1 A 4 a	3.78%			2 D 2	2.76%
	1 A 3 b i	0.92%	1 A 4 c i	3.56%			1 A 3 b v	2.62%
		1 A 4 c iii	3.50%			1 A 3 b iii	2.35%	
		1 A 1 b	2.65%			3 B	2.11%	
Energy		1 A 2 a	2.05%			1 A 3 b ii	1.95%	
Transport						1 A 3 d ii	1.91%	
Industry						1 B 2 a v	1.70%	
Solvent and product use						1 A 4 c ii	1.27%	
Agriculture						1 A 2 f	1.04%	
Waste						1 A 4 c i	0.86%	
Other								
Total (%)		95.47%		95.52%		95.20%		95.32%

Table 1.1.b. Key sources for CO and particulate matter species

CO		TSP		PM10		PM2.5	
1 A 3 b i	37.91%	2 G	15.45%	4 B 9	12.63%	1 A 3 b i	10.63%
1 A 2 a	10.87%	4 B 9	9.98%	2 G	8.56%	1 A 3 b iii	9.36%
2 C	9.53%	2 C	9.11%	4 B 8	6.17%	1 A 3 b ii	8.28%
1 A 4 b i	9.01%	1 A 4 b i	6.55%	1 A 3 b i	6.14%	1 A 4 b i	8.07%
1 A 3 b iv	8.88%	4 B 8	4.88%	2 C	5.65%	1 A 4 c ii	6.51%
1 A 2 c	3.97%	1 A 3 b i	4.86%	1 A 3 b iii	5.41%	1 A 3 b vi	6.42%
1 A 3 d ii	3.32%	2 B 5	4.31%	1 A 3 b ii	4.79%	7	6.30%
1 A 3 b iii	3.01%	1 A 3 b iii	4.28%	1 A 4 b i	4.67%	1 A 3 b vii	5.51%
1 A 3 b ii	2.56%	2 D 2	3.79%	1 B 2 a iv	4.22%	2 C	4.89%
1 A 2 f	2.05%	1 A 3 b ii	3.78%	1 A 4 c ii	3.76%	1 B 2 a iv	4.38%
1 A 4 c ii	2.00%	1 B 2 a iv	3.39%	1 A 3 b vi	3.71%	4 B 9	4.37%
1 A 1 a	1.38%	1 A 4 c ii	2.97%	7	3.64%	1 A 2 e	3.05%
1 A 2 e	0.85%	1 A 3 b vi	2.93%	2 D 2	3.39%	1 A 3 d ii	3.04%
		7	2.88%	1 A 3 b vii	3.18%	1 A 2 f	2.94%
		1 A 3 b vii	2.52%	2 A 7	3.06%	4 B 8	2.14%
		2 A 7	2.44%	1 A 2 e	2.94%	2 B 5	2.10%
		1 A 2 e	2.36%	3 D	2.91%	2 A 7	1.85%
		3 D	2.30%	1 A 2 f	2.83%	1 A 1 a	1.65%
		1 A 2 f	2.25%	2 B 5	2.21%	2 G	1.48%
		4 B 1 a	1.70%	4 B 1 a	2.15%	1 A 4 c iii	1.25%
		1 A 1 a	1.61%	1 A 3 d ii	1.76%	2 D 2	0.88%
		1 A 3 d ii	1.39%	1 A 1 a	1.19%		
				2 D 1	0.95%		
	95.34%		95.72%		95.92%		95.10%

Table 1.1.c. Key sources for Pb, Hg, Cd, DIOX and PAH

Pb		Hg		Cd		DIOX		PAH	
2 C	59.66%	1 A 1 a	34.18%	2 B 5	46.87%	3 D	64.37%	3 D	44.82%
1 A 3 b vi	15.69%	2 B 5	22.29%	2 C	43.08%	1 A 4 b i	15.09%	2 G	13.64%
2 A 7	10.32%	2 C	21.18%	1 A 4 b i	3.49%	1 A 2 a	6.62%	1 A 4 b i	13.36%
1 A 4 b i	5.71%	2 A 1	9.75%	1 A 1 a	2.92%	1 A 1 a	6.48%	1 A 3 b i	8.75%
2 B 5	5.37%	6 C	9.36%			1 A 3 b i	1.45%	1 A 3 b iii	6.55%
						6 C	0.89%	1 A 3 b ii	3.53%
						2 A 7	0.89%	1 A 2 f	2.83%
								1 A 3 b iv	1.45%
								1 A 4 c ii	1.37%
	96.75%		96.76%		96.37%		95.77%		96.30%

1.6 Reporting, QA/QC and archiving

Reporting

The Informative Inventory Report is prepared by MNP, with contributions from experts from the PER task forces. The Ministry of VROM formally approves the IIR before it is submitted, in some cases following consultation with other ministries.

QA/QC

MNP as well as the PER are ISO 9001:2000 certified. QA/QC, documentation and archiving is done according to procedures of the quality manual. Arrangements and procedures for the contributing institutes are described in the yearly project plan (MNP, 2006).

In general, the following QA/QC activities are performed:

- **QC checks.** A number of general QC checks have been introduced as part of the annual work plan of the PER. The QC checks built into the work plan aim at covering such issues as consistency, completeness and accuracy of the NFR data.

The general QC for the inventory is largely performed within the PER as an integrated part of the working processes. For the 2006 inventory the PER task forces filled in a standard-format database with emission data for 1990, 1995, 2000 and 2003–2005. After a first check of the emission files by MNP and TNO for completeness, the (corrected) data became available to the specific task force for checking consistency and trend analysis (comparability, accuracy). The task forces have access to information about the relevant emissions in the database. Several weeks before the data set is fixed, a trend verification workshop is organised by MNP (see Box 1.1).

Box 1.1. Trend verification workshops

Several weeks in advance of a trend analysis meeting, a snapshot from the database is made available by MNP in a web-based application (Emission Explorer, EmEx) for checks by the institutes involved and experts (PER task forces). In this way the task forces can check for level errors and consistency in the algorithm/method used for calculations throughout the time series. The task forces perform checks for relevant gases and sectors. The totals for the sectors are then compared with the previous year's data set. Where significant differences are found, the task forces evaluate the emission data in more detail. The results of these checks form the subject of discussion at the trend analysis workshop and are subsequently documented.

Furthermore, TNO provides the task forces with time series of emissions per substance for the individual target sectors. The task forces examine these time series. During the trend analysis for this inventory the emission data were checked in two ways: 1) emissions from 1990 – 2004 from the new time series were compared with the time series of last years inventory and 2) the data for 2005 were compared with the trend development per gas since 1990. The checks of outliers are performed on a more detailed level of the sub-sources of all sector background tables:

- annual changes in emissions;
- annual changes in activity data;
- annual changes in implied emission factors and
- level values of implied emission factors.

Exceptional trend changes and observed outliers are noted and discussed at the trend analysis workshop, resulting in an action list. Items on this list have to be processed within 2 weeks or be dealt with in next year's inventory.

Quality assurance (QA)

QA activities can be summarised as follows:

- For the energy, industry and waste sectors, emission calculation in the PER is based mainly on annual environmental reports by companies. The companies themselves are responsible for the data quality; the competent authorities (in the Netherlands, mainly provinces and local authorities) are responsible for checking and approving the reported data, as part of the annual quality assurance.
- As part of the evaluation process of the previous cycle, internal audits are performed within MNP as part of the ISO certification.
- Furthermore, QA checks are planned to be performed by institutes actually not involved in the PER system.

Archiving and documentation

Internal procedures are agreed on (for example, in the PER work plan) for general data collection and the storage of fixed data sets in the PER database at MNP, including the documentation/archiving of QC checks. Moreover, updating of monitoring protocols for substances under the Convention for Long Range Transboundary Air Pollution is one of the priorities within the PER system. Emphasis is put on documentation of methodologies for calculating SO_x, NO_x, NMVOC, NH₃ and PM₁₀ (PM_{2.5}). Methodologies/ protocols, emission data (including the emissions of Large Point Sources on the basis of annual Environmental Reports), as well as such emission reports as the National Inventory Report (UNFCCC) and the Informative Inventory Report (CLRTAP), are made available on the website of the PER: www.prtr.nl or www.emissieregistratie.nl (Dutch version). Each institution involved in the PER is responsible for QA/QC aspects related to reports based on the annually fixed database.

1.7 Uncertainties

Uncertainty assessments constitute a means to either provide the inventory users with a quantitative assessment of the inventory quality or to direct the inventory preparation team to priority areas, where improvements are warranted and can be made cost-effective. For these purposes, quantitative uncertainty assessments have been carried out for these purposes since 1999. However, awareness of uncertainties in emission figures was expressed earlier in the PER in so-called quality indices and in several studies on industrial emissions and generic emission factors for industrial processes and diffuse sources. To date, the Dutch PER gives only one value for emissions (calculation result, rounded off to three significant digits).

The information on the quality of emission figures presented here is based on the TNO report 'Uncertainty assessment of NO_x, SO₂ and NH₃ emissions in the Netherlands' (Van Gijlswijk et al., 2004), which presents the results of a Tier-2 'Monte Carlo' uncertainty assessment.

1.7.1 Quantitative uncertainty

Uncertainty estimates in national total emissions have been reported in the Environmental Balances since 2000 (RIVM, 2001). These estimates were based on uncertainties by source category using simple error propagation calculations (Tier 1). Most uncertainty estimates are based on the judgement

of RIVM/MNP emission experts. A preliminary analysis on NMVOC emissions showed an uncertainty range of about 25%. In a recent study by Van Gijlswijk et al. (2004), the uncertainty in the contribution of the various emission sources to total acidification (in acidification equivalents) was assessed according to the Tier-2 methodology (estimation of uncertainties by source category using Monte Carlo analysis). See Table 1.2 for results. A comparison was also made between the Tier-1 and Tier-2 methodologies. This is not straightforward as the two studies use a different knowledge collection. The 2000 Tier-1 analysis used CLRTAP default uncertainties for several NO_x processes, which explains the difference with the 1999 Tier-1 results. For NH₃, the difference between the 2000 Tier-1 and Tier-2 can be explained by taking non-normal distributions and dependencies between individual emission sources for each animal type into account (both are violations of the Tier-1 assumptions: effects encapsulated in the 1999 Tier-1 analysis). The differences for SO₂ and total acidifying equivalents are small. The conclusion drawn from this comparison is that focusing on the order of magnitude of the individual uncertainty estimates, as in the RIVM (2001) study, provides a reasonable first assessment of the uncertainty of source categories.

Table 1.2. Uncertainty (95% confidence ranges) in acidifying compounds and for total acidifying equivalents for emissions in 1999 (RIVM, 2001) and 2000 (Van Gijlswijk et al., 2004)

Component	Tier-1 for 1999	Tier-1 for 2000	Tier-2 for 2000
NH ₃	± 17%	± 12%	± 17%
NO _x	± 11%	± 14%	± 15%
SO ₂	± 8%	± 6%	± 6%
Total acid equivalents	± 9%	± 8%	± 10%

The RIVM (2001) study draws on the results of an earlier study on the quality of nitrogen oxides (NO_x) and sulphur dioxide (SO₂) emissions, as reported by individual companies for point sources under their national reporting requirements. Besides providing quantitative uncertainty estimates, the study yielded important conclusions. One example was that a limited number of facilities showed high uncertainties (e.g. 50% or more for NO_x), which could be reduced with little extra effort, and that companies generally have a lack of knowledge on the uncertainty about the emissions they report.

In the study by Van Gijlswijk (2004), emission experts were systematically interviewed on quantitative uncertainties, which provided simultaneous information on the reliability and quality of the underlying knowledge base. For processes not covered by interviews, standard default uncertainties, derived from the Good Practice Guidance for CLRTAP emission inventories, were used (Pulles and Van Aardenne, 2001). The qualitative knowledge (on data validation, methodological aspects, empirical basis and proximity of data used) has been combined into a score for data strength, based on the so-called NUSAP approach (Van der Sluijs et al., 2003; Van der Sluijs et al., 2005). The qualitative and quantitative uncertainties were combined in so-called diagnostic diagrams that can be used to identify areas for improvement, since the diagrams indicate strong and weak parts of the available knowledge (see Figure 4.1). Sources with a relatively high quantitative uncertainty and weak data strength are thus candidates for improvement. To effectively reduce the uncertainty, the nature of uncertainties must be known (e.g. random, systematic or knowledge uncertainty). A general classification scheme on uncertainty typology is given in Van Asselt (2000).




Figure 1.2. NUSAP diagnostic diagram indicating strong and weak elements in the available knowledge on acidifying substances.

1.8 Explanation on the use of notation keys

The Dutch emission inventory covers all relevant sources as specified in the CLRTAP that determine the emissions to air in the Netherlands. Because of the long history of the inventory it is not always possible to specify all subsectors in detail. This is the reason notation keys are used in the emission tables (NFR). These notation keys will be explained in tables 1.3 – 1.5.

Table 1.3 The NE notation key explained

NFR code	Substance(s)	Reason for reporting NE
All	Benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene, Indeno(1,2,3-cd)pyrene	Currently, the inventory does not provide data on individual compounds. These are included in the category Total 1-4, which is filled with total PAH emissions according to the Dutch definition of the 'VROM sum of 10 PAH'.

Table 1.4. The IE notation key explained

NFR code	Substance(s)	Included in NFR code
1 A 3 a ii (ii)	All	No specific data are available on the domestic (very small) cruise emissions. These emissions are incorporated in 1 A 3 a ii (i) (based on total fuel use for domestic flights).
1 B 2 c	All	Venting and flaring emissions occur almost exclusively in the natural gas sector and are therefore included in 1 B 2 b.
4 B 1 b	TSP, PM ₁₀ , PM _{2.5}	Since no specific data are available for this subcategory, all emissions are reported under 4 B 1 a.

Table 1.5. Sub-sources accounted for in reporting 'other' codes

NFR code	Sub-source description	Substance(s) reported
1A2f	Combustion in the non-specified industries, machineries, services, production activities.	All
1A3 e	NO/NA	
1A5a	Combustion of landfill gas	All
1A5b	NO/NA	
1B1 c	NO/NA	
1B2 a vi	NO/NA	
2 A 7	Process emissions in construction activities and production of building materials	All
2 B 5	Process emissions during production of chemicals, paint, pharmaceuticals, soap, detergents, glues and other chemical products	All
2 G	Process emissions during production of wood products, plastics, rubber, metal, textiles and paper	All
3 D	Use of products, not in 3A-C, venting transport and storage facilities; use of products by consumers, in commercial activities, tobacco products for smoking (NMVOC only), cooling, freezing and air-conditioning	All
4 B 13	Pets	NH ₃
4 G	Handling agricultural-based materials and products	NH ₃ , TSP, PM ₁₀ , PM _{2.5}
6 D	Handling waste	all
7	Tobacco products for smoking Transpiration, breathing	All substances, excl NMVOC NH ₃
5E	NO/NA	

1.9 Missing sources

The Netherlands emission inventory covers all important sources. However, no data is available for individual PAHs. The reason for this is the less restrictive requirements on the individual PAHs to be reported in environmental reports from companies, in such a way that it is not possible to speciate the total PAH emissions. A study performed on priority substances (PAH emissions; heavy metals; (other) Persistent Organic Pollutants) in 2005 included a number of recommendations for improving reporting speciated PAHs (Alkemade et al., 2005). These recommendations are further elaborated in 2007/2008, aiming at providing individual PAH data in the next submission of the IIR.

2 Trends in emissions

2.1 Trends in national emissions

The emissions of all gases showed a downward trend in the 1990 – 2005 period (see Table 2.1). The major overall drivers for this trend are:

- emission reductions in the industrial sectors,
- cleaner fuels and
- cleaner cars.

European regulations for road traffic have caused a decrease in vehicle emissions of 72% since 1990 for NMVOC, 47% for particulate matter, 45% for NO_x and about 92% for SO₂, despite a growth in traffic of 33%. For particulate matter and NO_x, standards have been set for installations by tightening up the extent of emission stocks of heating installations (BEES). In meeting these requirements Dutch industrial plants have realised a reduction of 70% in particulate matter emissions and 50% in NO_x emissions since 1990. The drivers for the downward emission trend for specific gases will be elaborated in more detail in the next section.

Table 2.1. Total national emissions, 1990-2005

Year	Main Pollutants					Particulate Matter			Priority Heavy Metals		
	NO _x	CO	NMVOC	SO _x	NI13	TSP	PM10	PM2.5	Pb	Cd	Hg
	Gg	Gg	Gg	Gg	Gg	Gg	Gg	Gg	Mg	Mg	Mg
1990	558	1136	465	190	250	100	76	46	340	2,1	4,7
1995	468	860	332	128	193	74	57	34	164	1,1	1,2
2000	394	721	234	72	152	57	46	27	35	1,0	0,9
2003	373	624	187	63	135	54	41	24	40	2,4	0,7
2004	355	615	180	64	134	51	40	23	44	1,8	1,0
2005	344	599	176	62	135	50	40	23	44	1,7	1,0
period 1990-2005, abs	-214	-537	-289	-128	-114	-50	-37	-23	-296	-0,4	-3,6
period 1990-2005, %1990	-38%	-47%	-62%	-67%	-46%	-50%	-48%	-50%	-87%	-18%	-78%

Year	Persistent Organic Pollutants (POPs) Annex I (1)									POPs Annex II (2)		
	Alidin	Chlordane	Chlordane	Dieldrin	Endrin	Heptachlor	Hexabromo-biphenyl	Mirex	Toxaphene	HCH	DDT	PCB
	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg	kg
1990	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
1995	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2000	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2003	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2004	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
2005	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
period 1990-2005, abs												
period 1990-2005, %1990												

Year	POPs Annex III (3)						Other Heavy Metals						Other POPs (4)		
	DiOX	PAH					HCB	As	Cr	Cu	Ni	Se	Zn	PCP	SCCP
g I-Teq	benz(a)pyrene	benz(b)fluoranthene	benz(k)fluoranthene	indene(1,2,3-cd)pyrene	Total 1-4	Mg									
1990	743	NO	NO	NO	NO	1710	0	1,5	10,0	70,1	75,3	0,4	225,6	34000	NO
1995	66	NO	NO	NO	NO	900	0	1,0	6,7	74,8	86,6	0,3	148,6	29000	NO
2000	31	NO	NO	NO	NO	568	0	1,1	3,2	82,4	19,4	0,5	100,1	24000	NO
2003	26	NO	NO	NO	NO	429	0	1,0	2,9	86,7	12,6	0,7	96,8	22050	NO
2004	28	NO	NO	NO	NO	490	0	1,5	3,2	87,4	13,8	1,1	93,1	21400	NO
2005	28	NO	NO	NO	NO	483	0	1,5	3,2	87,9	13,4	1,1	93,1	21400	NO
period 1990-2005, abs	-715					-1228	0	0,0	-6,8	17,8	-61,9	0,7	-133	-12600	
period 1990-2005, %1990	-96%					-72%		3%	-68%	25%	-82%	177%	-59%	-37%	

Notes: PAH emissions should be 'NE' and HCB emissions should be 'NO'.

2.2 Trends for sulphur dioxide (SO₂)

The Dutch SO_x emissions (reported as SO₂) decreased by 128 Gg in the 1990 – 2005 period, corresponding to 67% of the national total in 1990 (Figure 2.1). Main contributions to this decrease came from the energy, industry and transport sectors. The use of coal declined and major coal-fired electricity producers installed flue gas desulphurisation plants. The sulphur content in fuels for the (chemical) industry and traffic was also reduced. Currently, the energy sector is responsible for almost three-quarters of the national SO₂ emission. The traffic share has grown because of the increase in diesel-powered vehicles.




Figure 2.1. SO₂, emission trend 1990-2005 and share by sector in 1990 and 2005

2.3 Trends for nitrogen oxides (NO_x)

The Dutch NO_x emissions (NO and NO₂, expressed as NO₂) decreased by 214 Gg in the 1990 to 2005 period; this corresponded to 38% of the national total in 1990 (Figure 2.2). Main contributors to this decrease were the road transport and energy sectors. The emissions per vehicle decreased significantly in this period, but were counterbalanced by an increase in number and mileages of vehicles. The share of the different NFR categories in the national total did not change significantly.




Figure 2.2. NO_x emission trend 1990-2005 and share by sector in 1990 and 2005

2.4 Trends for ammonia (NH₃)

The Dutch NH₃ emissions decreased by 114 Gg in the 1990 to 2005 period, corresponding to 46 % of the national total in 1990 (Figure 2.3). This decrease was due to the agricultural sources. The direct emissions from animal husbandry increased but measures were taken to reduce the emissions during application of manure to the soil. Now, over 90 % of the Dutch NH₃ emissions come from agricultural sources.




Figure 2.3. NH₃, emission trend 1990-2005 and share by sector in 1990 and 2005

2.5 Trends for non-methane volatile organic compounds (NMVOC)

The Dutch NMVOC emissions decreased by 289 Gg in the 1990 to 2005 period, corresponding to 62% of the national total in 1990 (Figure 2.4). All major source categories contributed to this decrease, for example transport (introduction of catalyst and cleaner engines), product use (intensive programme to reduce NMVOC content in consumer products and paints) and industry (introducing emission abatement specific for NMVOC).




Figure 2.4. NMVOC, emission trend for 1990-2005 and share by sector in 1990 and 2005

2.6 Trends for particulate matter (PM₁₀)

The Dutch PM₁₀ emissions decreased by 37 Gg in the 1990 to 2005 period, corresponding with 48% of the national total in 1990 (Figure 2.5). The major source categories contributing to this decrease are:

- industry (combustion and process emissions) due to cleaner fuels in refineries and the side-effect of emission abatement for SO₂ and NO_x ;
- traffic and transport, and
- introduction of cleaner diesel engines.




Figure 2.5. PM₁₀ emission trend 1990-2004 and share by sector in 1990 and 2004

The emissions from animal husbandry in agriculture did not change significantly; neither did the emissions from consumers (1A4b1). The share of the emissions from residential wood stoves increased from 42% for 1A4 in 1990 to 45% in 2005. $PM_{2.5}$ emissions are also included in the 2006 submission to UNECE. These emissions are calculated as a specific fraction of PM_{10} by sector (based on Wesselink et al., 1998). About 74% of the $PM_{2.5}$ emissions stem from combustion processes, especially from mobile combustion (road transport).

2.7 Trends for heavy metals (Pb and Cd)

The Dutch lead (Pb) emissions decreased by 296 Gg in the 1990 to 2005 period, corresponding to 87% of the national total in 1990 (Figure 2.6). This decrease is solely attributable to the transport sector, where, due to the removal of Pb from petrol, the Pb emissions collapsed. The remaining sources for Pb are the iron and steel industry, and industry (combustion and process emissions).




Figure 2.6. Pb, emission trend 1990-2005 and share by sector in 1990 and 2005




Figure 2.7. Cd, emission trend 1990-2005 and share by sector in 1990 and 2005

The Dutch cadmium (Cd) emissions decreased by nearly 400 kg in the 1990 to 2005 period, corresponding to 18 % of the national total in 1990 (Figure 2.7). This decrease is caused mainly by the large decrease in the emissions from waste combustion. Between 1990 and 2005 old incinerators without flue gas cleaning were closed, and state of the art emission abatement was installed in the both the remaining incinerators and, sometimes, in the newly built ones. The remaining major sources for Cd emissions in the Netherlands are the chemical industry and the iron and steel industry.

2.8 Trends for polycyclic aromatic hydrocarbons (PAH) and dioxins

The Dutch PAH emissions decreased by 1.2 Gg in the 1990 to 2005 period, corresponding to 72% of the national total in 1990 (Figure 2.8). Please note that the Netherlands reports its PAH emissions according to a specific Dutch definition, namely, the 'sum 10 PAH of VROM'. This definition not only includes the four PAH substances stated in NFR, but also six others. The Dutch PAH emission is therefore by definition higher than that given in the NFR definition. The major contributors to this decrease are the:

- (metal) industry (general emission reduction) and
- product use (ban on creosoted wood in several applications).

Please note that the Netherlands reports only the total PAH emission. The recommended detailed speciation is not yet available. Further actions to derive the detailed figures from the individual PAH from the PAH total will be necessary, see section 1.9.




Figure 2.8. PAH, emission trend 1990-2005 and share by sector in 1990 and 2005




Figure 2.9. Dioxins, emission trend in 1990-2005 and share by sector in 1990 and 2005

Dioxin emissions decreased by 715 g I-Teq in the 1990 to 2005 period, corresponding to 96% of the national total in 1990 (Figure 2.9). In the period after 1990 specific emission abatement introduced into all waste incineration plants, was specifically targeted to reduce dioxin emissions. Furthermore, measures were taken to reduce dioxin emissions in the energy and industrial sectors. Currently, the major source for dioxin emission is the category product use (3).

3 Energy, stationary fuel combustion (1A)

About 80-100% of the NO_x, SO₂, PM₁₀ and NH₃ emissions from stationary combustion (categories 1A1, 1A2, 1A4 and 1A5) are based on environmental reports of large industrial companies. The emission data in the Environmental Annual Reports (EARs) are based on direct emission measurements (see formula below) or calculations based on fuel input and emission factors. The emission factors used in the calculations are also based on measurements according to this formula.

[Concentration]*Flow*Duration of emission

where:

[Concentration] = Online (semi-)continuous measurement: frequency - seconds to daily

Discontinuous measurement: several times a year, directly in air flow

Off line: sampling and analysis in laboratory

Flow = Flow-speed measurement in air flow; surface of flow channel; calculation based on fuel or raw materials/production quantities; At diffuse emissions: calculation or air flow over source

Duration = Calculation based on process-control data

The emissions and fuel consumption data in the EARs are systematically examined for inaccuracies by checking the resulting implied emission factors. If the environmental report provides data of high enough quality (see section 1.3 on QA/QC), the information is used to calculate an 'implied emission factor' for a cluster of reporting companies (aggregated by SBI code) and the Emission factor ER-I. These emission factors are fuel and sector dependent.

Emission factor ER-I (SBI category, fuel type) =

Emissions ER-I (SBI category, fuel type) / Energy use ER-I (SBI category, fuel type)

Where:

ER-I = Emission Registration database for individual companies

Next, the total combustion emissions in this SBI category are calculated from the energy use, as provided in the Netherlands Energy Statistics (Statistics Netherlands), multiplied by the implied emission factor.

ERI_SBI_Emission (SBI category, fuel type) =

Emission factor ER-I (SBI category, fuel type) * Energy use NEH (SBI category, fuel type)

For sectors with no individual registration of emissions (e.g. residential and agricultural sectors), a set of specific emission factors is used (see section 3.3).

3.1 Energy industries (1A1)

Public electricity and heat production (1A1a)

Emission data are based on environmental annual reports and collectively estimated industrial sources. For this source category, the percentages of emissions based on annual reports are: 90% for NO_x, 80% for SO₂, 90% for CO and 100% for Hg, Cd and dioxins.

Category 1A1a is a key source for the following components (% of national total in 2005):

SO _x	(17.1%)
NO _x	(12.8%)
CO	(1.4%)
TSP	(1.6%)
PM ₁₀	(1.2%)
PM _{2.5}	(1.7%)
Hg	(34.2%)
Cd	(2.9%)
DIOX	(6.5%)

Petroleum refining (1A1b)

All emission data are based on Environmental Annual reports and registered in the ER-I database.

Category 1A1b is a key source for the following components (% of national total):

SO _x	(47.3%)
NO _x	(2.7%)

Manufacture of solid fuels and other energy industries (1A1c)

No key sources in this category.

3.2 Manufacturing industries and construction (1A2)

Iron and steel (1A2a)

All emission data are based on environmental annual reports and registered in the ER-I database.

Category 1A2a is a key source for the following components (% of national total):

SO _x	(6.4%)
NO _x	(2.0%)
CO	(10.9%)
DIOX	(6.6%)

Non-ferrous metals (1A2b)

Emission data are based on environmental annual reports and collectively estimated industrial sources. For this source category, the percentage of SO₂-emissions based on annual reports is 100%.

Category 1A2b is a key source for the following components (% of national total):

SO _x	(5.3%)
-----------------	--------

Chemicals (1A2c)

Emission data are based on Environmental Annual reports and collectively estimated industrial sources. For this source category, the percentages of emissions based on annual reports are about 100% for SO₂, 90% for NO_x, 75% for CO and 100% for Pb, Cd and dioxins.

Category 1A2c is a key source for the following components (% of national total):

SO_x	(6.5%)
NO_x	(4.3%)
CO	(4.0%)

Pulp, paper and print (1A2d)

All emission data are based on Environmental Annual reports and registered in the ER-I database. No key sources are found in this category.

Food processing, beverages and tobacco (1A2e)

Emission data are based on Environmental Annual reports and collectively estimated industrial sources. No key sources are found in this category.

CO	(0.9%)
TSP	(2.4%)
PM₁₀	(2.9%)
PM_{2.5}	(3.1%)

Other (1A2f)

This sector includes all combustion emissions from the industrial sectors not belonging to the categories 1A2a to 1A2f. Emission data are based on Environmental Annual reports and collectively estimated industrial sources.

Category 1A2f is a key source for the following components (% of national total):

SO_x	(5.4%)
NO_x	(6.0%)
NMVOC	(1.0%)
CO	(2.1%)
TSP	(2.3%)
PM₁₀	(2.8%)
PM_{2.5}	(2.9%)
PAH	(2.8%)

3.3 Other sectors (1A4)**Commercial / institutional (1A4a)**

Combustion emissions from the commercial and institutional sector are based on fuel consumption data (Statistics Netherlands) and emission factors (see Table 3.1.).

Table 3.1. Emission factors for stationary combustion emissions of the services sector (g/GJ)

	Natural gas	Domestic fuel oil	LPG	Paraffin oil	Coal	Oil fuel
VOC	30	10	2	10	35	10
SO ₂	0.22	87	0.22	4.6	460	450
NO _x	1)	50	40	50	300	125
CO	10	10	10	10	100	10
Carbon black		5	10	2		50
Fly ash					100	
PM ₁₀	0.15	4.5	2	1.8	2	45
PM coarse		0.5		0.2	80	5

1) see table on NO_x emission factors in Soest-Vercammen et al. (2002)

Category 1A4a is a key source for the following components (% of national total):

NO _x	(3.8%)
-----------------	--------

Residential (1A4b)

1A4bi Residential plants

Combustion emissions of central heating, hot water and cooking are based on fuel consumption data (Statistics Netherlands) and emission factors (see Table 3.2.). The major fuel used in this category is natural gas. The use of wood in stoves and fireplaces for heating is almost negligible.

Combustion emissions of (wood) stoves and fireplaces are calculated by multiplying the fuel consumption by apparatus type and by fuel type (Statistics Netherlands) with emission factors per house (Hulskotte et al., 1999).

Table 3.2. Emission factors for combustion emissions of households (g/GJ)

	Natural gas	Domestic fuel oil	LPG	Paraffin oil	Coal
VOC	6.3	15	2	10	60
SO ₂	0.22	87	0.22	4.6	420
NO _x	1)	50	40	50	75
CO	15.8	60	10	10	1500
Carbon black	0.3	5	10	2	
Fly ash					200
PM ₁₀	0.3	4.5	2	1.8	120
PM coarse		0.5		0.2	80

1) See table on NO_x emission factors in Soest-Vercammen et al. (2002)

Category 1A4b i is a key source for the following components (% of national total):

NO_x	(5.5%)
NMVO	(4.9%)
CO	(9.0%)
TSP	(6.6%)
PM₁₀	(4.6%)
PM_{2.5}	(8.1%)
Pb	(5.7%)
Cd	(3.5%)
DIOX	(15.1%)
PAH	(13.4%)

1A4bii Household and gardening (mobile)

Emissions are included in category 1A4b i and can not be separated due to lack of specific fuel data on this level.

Agriculture / forestry / fishing (1A4c)

1A4ci Stationary

Stationary combustion emissions are based on fuel consumption obtained from Statistics Netherlands, which is, in turn, based on data from the Agricultural Economics Research Institute, and emission factors (Table 3.3).

Table 3.3. Emission factors for stationary combustion emissions from agriculture (g/GJ)

	Natural gas	Domestic fuel oil	LPG	Paraffin oil	Coal	Oil fuel
VOC	30	10	2	10	35	10
SO ₂	0.22	87	0.22	4.6	460	450
NO _x	1)	50	40	50	300	125
CO	10	10	10	10	100	10
Carbon black		5	10	2		50
Fly ash					100	
PM ₁₀	0.15	4.5	2	1.8	2	45
PM coarse		0.5		0.2	80	5

1) See table on NO_x emission factors in Soest-Vercammen et al. (2002)

Category 1A4ci is a key source for the following components (% of national total):

NO_x	(3.6%)
NMVO	(0.9%)

1A4cii Off-road vehicles and other machinery

Combustion emissions of CO, VOC, NO_x, PM₁₀, SO₂ and heavy metals from off-road vehicles and other machinery are based on diesel fuel consumption and emission factors (g/kg fuel). Fuel consumption data from private farm machinery is provided by the Agricultural Economics Research

Institute LEI, while data for agricultural work contractors is provided by Statistics Netherlands. Fuel consumption in the building sector is based on production statistics of this sector, provided by Statistics Netherlands. The fuel consumption of other machinery is highly uncertain, as it is based on the difference between the total amount of diesel oil used according to the Netherlands Energy Statistics minus the use of diesel oil in the agricultural and construction sector. Combustion emissions of NH₃ are based on EEA emission factors (Ntziachristos and Samaras, 2000) and total fuel consumption by off-road vehicles and other machinery. VOC and PAH combustion emissions are calculated using VOC profiles (VROM, 1993 and Shareef et al., 1988).

Category 1A4cii is a key source for the following components (% of national total):

SO_x	(2.5%)
NO_x	(5.5%)
NMVOG	(1.3%)
CO	(2.0%)
PM₁₀	(3.8%)
PM_{2.5}	(6.5%)
PAH	(1.4%)

1A4ciii National fishing

Combustion emissions are based on fuel sales to cutters operating within national waters and fuel specific emission factors. Since fuel sales to cutters are not recorded separately in the Netherlands Energy Statistics (these are contained in the bunker fuel sales) an estimate of fuel use is made on the basis of ship movements. Emission factors for CO, NO_x, (NM)VOC, CH₄, SO₂, and PM₁₀ are derived from national research (Hulskotte and Koch, 2000; Van der Tak, 2000). NH₃ emission factors are derived from Ntziachristos and Samaras (2000). It is assumed that all four-stroke engines use diesel oil, while all two-stroke engines use heavy fuel oil. VOC and PAH combustion emissions are calculated using VOC profiles (VROM, 1993 and Shareef et al., 1988).

Category 1A4ciii is a key source for the following components (% of national total):

SO_x	(1.1%)
-----------------------	--------

3.4 Other (1A5)

Other, stationary (including military) (1A5a)

Emissions in this category are wrongly reported in category 1A5b. This will be corrected in the next submission.

Other, mobile (including military) (1A5b)

For military vessels and aircraft only emissions of CO₂, N₂O and CH₄ are calculated. Other compounds relating to NEC ceilings can not be calculated, since it is unknown where fuel is used. The Ministry of Defence regards information on the location of military activity as classified.

No key sources are found in this category.

3.5 Mobile combustion (1A3)

Road transportation (1A3b)

Exhaust emissions of CO, NMVOC, NO_x, NH₃ and PM₁₀ in these source categories are dependent on fuel type, emission reduction technology, and vehicle type and vehicle use. These emissions are calculated on the basis of traffic performance (vehicle kilometres) and specific emission factors for a variation of different vehicle classes and for three different road types. The vehicle classes are defined by the vehicle category (passenger car, van, etc.), fuel type, weight class, environmental class and in some instances the engine type and/or the emission reduction technology. The emission factors for passenger cars are based on a yearly monitoring programme by TNO (Built Environment and Geosciences). The specific emission factors per vehicle class are aggregated to emission factors by year of construction (in grams per vehicle kilometre). The emission factors by year of construction are published in [Statline](#), the central database of Statistics Netherlands. The method is described in detail in Klein et al. (2006).

Traffic performance data is based on the following data by Statistics Netherlands: ‘Survey on movement behaviour’, ‘Statistics on road freight transport’, ‘Motor cycling statistics’ (based on a survey in 1993) and ‘Mobility of Dutch residents’. The characteristics of the Dutch vehicle fleet are based on ‘Statistics on motor vehicles’, which in turn is based on data provided by the Dutch road traffic department (RDW). Passenger car movements by non-residents are based on the following data by Statistics Netherlands: ‘Statistics on registered overnight stays’, commuter traffic by foreign workers and number of day trips. Foreign freight transport kilometres are based on ‘Statistics on road freight transport’ and similar statistics from other EU countries provided by Eurostat.

Emissions of SO₂ and heavy metals (and CO₂) are dependent on fuel consumption and fuel type. These emissions are calculated by multiplying fuel use with emission factors (gram per litre fuel consumption). The emission factors are based on the sulphur, carbon and heavy metal contents of the fuels. It is assumed that 75% of the lead is emitted as particles and 95% of the sulphur is transformed to sulphur dioxide. The data on fuel consumption of mobile sources is collected by Statistics Netherlands.

Emissions of VOC components (alkanes, alkenes, aromates, such as benzene and formaldehyde, polycyclic aromatic hydrocarbons PAHs and chlorinated hydrocarbons) are calculated by multiplying the total VOC emission by a VOC speciation profile.

1A3bi Road transport, passenger cars

Category 13bi is a key source for the following components (% of national total):

SO ₂	(0.9%)
NO _x	(12.8%)
NH ₃	(1.8%)
NMVOC	(14.6%)

CO	(37.9%)
TSP	(4.9%)
PM₁₀	(6.1%)
PM_{2.5}	(10.6%)
DIOX	(1.5%)
PAH	(8.8%)

1A3bii Road transport, light duty vehicles

Category 1A3bii is a key source for the following components (% of national total):

NO_x	(5.5%)
NMVO	(2.0%)
CO	(2.6%)
TSP	(3.8%)
PM₁₀	(4.8%)
PM_{2.5}	(8.3%)
PAH	(3.5%)

1A3biii Road transport, heavy duty vehicles

Category 1A3biii is a key source for the following components (% of national total):

NO_x	(23.5%)
NMVO	(2.4%)
CO	(3.0%)
TSP	(4.3%)
PM₁₀	(5.4%)
PM_{2.5}	(9.4%)
PAH	(6.6%)

1A3biv Road transport, mopeds and motorcycles

Category 1A3biv is a key source for the following components (% of national total):

NMVO	(5.5%)
CO	(8.9%)
PAH	(1.5%)

1A3a Civil aviation

Combustion emissions – Amsterdam Airport Schiphol

Combustion emissions of CO, VOC, NO_x, PM₁₀, SO₂ and heavy metals from aviation are calculated with the EMASA model on a yearly basis (TNO Built Environment and Geosciences). This model is consistent with the US Environmental Protection Agency model for aviation emissions.

The Landing and Take-off cycle (LTO) can be divided into four LTO cycle phases: idle, take-off, climb-out and approach from 3000 feet. The four modes in the LTO cycle correspond to different power settings of the jet engines: idle 7%, take-off 100%, climb-out 85% and approach 30%. The equation for calculating the emissions is presented next:

$$\text{emission} = \text{SUM}_{p,m,f} (\text{LTO}_{p,m} * N_p * \text{FUEL}_{m,f} * \text{TIM}_{p,f} * \text{EF}_{m,f})$$

where:

emission = emission (kg/yr)

$\text{LTO}_{p,m}$ = number of LTO cycles per aircraft with jet engine type (m) per year

N_p = number of engines per aircraft

$\text{FUEL}_{m,f}$ = fuel consumption of jet engine type (m) in LTO cycle phase (f)

$\text{TIM}_{p,f}$ = time in mode in LTO cycle (f) for aircraft (p)

$\text{EF}_{m,f}$ = emission factor of jet engine type (m) in LTO cycle (f) (kg/kg)

The EMASA model takes into account about 100 types of aircrafts, as reported in the Statistical Annual Review of Amsterdam Airport Schiphol. The jet engine types of these aircrafts are based on the aircraft/engine combinations of the so-called Home-carriers (e.g. KLM, Martinair and Transavia). The emission factors originate in the DERA database (DERA, 1999), the Federal Aviation Agency Engine Emission Database (FAA, 1996); for smaller engines these are based on EPA publication AP42 (EPA, 1985). Emissions from military use of aviation fuel are reported under Other mobile sources (NFR 1A5b).

Emissions from auxiliary power units and general power units for aircraft at Schiphol are based on an estimated fuel consumption of 500 gram per passenger multiplied with emission factors.

Combustion emissions other airports

Emissions are calculated similarly to the method described above, now taking into account the number of flights per regional airport. Due to lack of data, splitting the flights according to aircraft type is done by the emission expert. Furthermore, emissions in the period 1995-1999 are calculated by indexing the 1994 emissions with the flights per airport in this period.

NH3 emissions are based on emission factors from EEA (Ntziachristos and Samaras, 2000) and total fuel consumption during the LTO cycle at Dutch airports.

VOC and PAH combustion emissions

First, the VOC emissions are calculated as described above. Second, the VOC and PAH components are calculated using VOC profiles (VROM, 1993 and Shareef et al., 1988).

1A3a(ii) Civil aviation (domestic, LTO)

No key source categories are found in this category.

1A3a(ii) Civil aviation (domestic, cruise)

Emissions are included in 1A3a(ii) and cannot be separated due to missing fuel data at this level.

Railways (1A3c)

Combustion emissions of CO, VOC, NO_x, PM₁₀, SO₂ and heavy metals from railways are based on diesel fuel consumption and emission factors. Fuel consumption data is provided by NS Reizigers

(Dutch rail passenger organisation). Combustion emissions of NH₃ are based on EEA emission factors (Ntziachristos and Samaras, 2000). VOC and PAH combustion emissions are calculated using VOC profiles (VROM, 1993 and Shareef et al., 1988).

No key sources are found in this category.

National navigation (1A3dii)

For inland ships energy consumption is calculated with a resistance model that is specific for the combination of waterway, boat type, direction of navigation and ship loading (Hulskotte et al., 2003). Emission factors dependent on energy consumption were derived by Oonk et al. (2003). Emission factors are dependent on year that ship was built and on maximum RPM for recently built engines. Energy consumption data is calculated using ship movements and divided into inland shipping and international shipping using the data of Statistics Netherlands.

Combustion emissions of leisure boats are based on fuel consumption data, which are estimated by multiplying boat numbers by specific yearly fuel consumption per boat type. Specific fuel consumption was determined by means of a questionnaire. The calculation procedure is described in a fact sheet (Hulskotte et al., 2005). Some of the emissions of some substances (e.g. PAH and NMVOC species) are specified as waterborne emissions.

Category 1A3dii is a key source for the following components (% of national total):

SO_x	(1.5%)
NO_x	(4.5%)
NMVOC	(1.9%)
CO	(3.3%)
TSP	(1.4%)
PM₁₀	(1.8%)
PM_{2.5}	(3.0%)

Other (1A3e)

No emissions are reported in this category and the subcategories 1A3ei *Pipeline compressors* and 1A3eii *Other mobile sources and machinery*.

3.6 Evaporation, tyre and brake wear, road abrasion (1A3b)

Road transport, gasoline evaporation (1A3bv)

VOC emissions from gasoline evaporation originate from diurnal losses, hot soak losses and running losses. The emission factors for gasoline evaporation are based on formulas from the CORINAIR project (Eggleston, 1989) and reported in Klein (1992). Since 1993, cars have been equipped with a charcoal canister in the fuel system. This has reduced the VOC emissions by 80%. For Euro 3 and Euro 4 vehicles, the reduction is assumed to be 90%. The emissions of VOC components are calculated on the basis of VOC speciation profiles. The evaporation VOC profile has, since 2000, been adjusted for the change in benzene and aromatics content of gasoline since 2000, due to stricter EU legislation (see Table 3.4).

Table 3.4. VOC profile for gasoline and gasoline evaporation (mass %)

	Gasoline		Gasoline vapour	
	1999 and before	2000 and after	1999 and before	2000 and after
Benzene	2.5	0.8	1	0.3
Toluene	15	12.5	3	2.5
Xylene	-	-	0.5	0.5
Aliphatic hydrocarbons (non-halogenated)	35	60	95	97
Aromatic hydrocarbons (non-halogenated)	65	40	5	3

Category 1A3bv is a key source for the following components (% of national total):

NMVO	(2.6%)
-------------	--------

Road transport, automobile tyre and brake wear (1A3bvi)

Particulate matter emissions (TSP) from tyre wear and brake wear are based on vehicle kilometres and emission factors. The fraction PM_{10} in total particulate matter for tyre wear is assumed to be 5% (highly uncertain) and for brake wear 100%. Heavy metal emissions are calculated using a speciation profile on total particulate emissions.

Category 1A3bvi is a key source for the following components (% of national total):

TSP	(2.9%)
PM_{10}	(3.7%)
$PM_{2.5}$	(6.4%)
Pb	(15.7%)

Road transport, automobile road abrasion (1A3bvii)

The same method is applied as for category 1A3bvi Tyre and brake wear. The fraction PM_{10} in total particulate matter for road abrasion is assumed to be 5% Heavy metal emissions are calculated using a speciation profile on total particulate emissions.

Category 1A3bvii is a key source for the following components (% of national total):

TSP	(2.5%)
PM_{10}	(3.2%)
$PM_{2.5}$	(5.5%)

3.7 Energy, fugitive emissions from fuels (1B)

The fugitive NMVOC emissions from category 1B2b comprise non-fuel combustion emissions from oil and gas production, emissions from gas transport (compressor stations) and gas distribution networks (pipelines for local transport).

The NMVOC emissions from oil and gas production and gas transport are derived from the environmental reports of the companies, which cover 100% of the emissions. The NMVOC emissions from gas distribution are calculated on the basis of a VOC profile with the CH₄ emission from the yearly report of the sector as input.

Category 1B2b is a key source for the following components (% of national total):

NMVOC	(3.7%)
-------	--------

4 Industry (2)

Industrial process emissions are based either on environmental reports of large industries or extrapolations to total emissions per SBI category, using implied emission factors and production data (method 1), or on sectoral reports on emissions (method 2), or specific emission factors and production statistics (CBS and trade organisations) (method 3).

Method 1 Extrapolation from emission data of individual companies

$$\text{Emission factor ER-I}_{(SBI\ category)} = \text{Emissions ER-I}_{(SBI\ category)} / \text{Production ER-I}_{(SBI\ category)}$$

where

ER-I = Emission Registration database for individual companies

Production ER-I = activity data or proxy for the production process

Next, the total process emissions in this SBI category are calculated from the production data, as provided in the Production Statistics (Statistics Netherlands), multiplied by the implied emission factor.

$$\text{ERI_SBI_Emission}_{(SBI\ category)} = \text{Emission factor ER-I}_{(SBI\ category)} * \text{Production}_{(SBI\ category)}$$

Note: Companies do not provide specific information to the PER on their measurement systems or emission model or which emission factors are used in the calculation model. Therefore, in some cases the PER can not use the data from the environmental reports in the extrapolation to the total emissions of a sector.

Method 2 Sectoral emission reports

Some trade organisations provide (yearly) emission reports as part of their agreements in covenants with the government; see <http://www.fo-industrie.nl> (Dutch only). Emissions reported by individual companies are subtracted from the total emissions reported by the trade organisation.

Method 3 Sectors with no individual registration

A set of specific emission factors is used for sectors with no individual registration of emissions, mostly based on the so-called SPIN documents, the ‘Cooperation project on industrial emissions’.

In this project the RIVM, assisted by consultant firms, revised and extended the original material (individual registration of about 6000 companies collected by TNO between 1974 and 1983); they also added proposals for abatement methods. These reports document about 90 industrial processes in the Dutch industry. The emission factors are combined with production statistics from CBS or activity data reported by specific trade organisations.

4.1 Mineral production (2A)

This category comprises emissions related to the production and use of non-metallic minerals in:

- 2A1 Cement clinker production;
- 2A3 Limestone and dolomite use;
- 2A4 Soda ash production and use;
- 2A7 Other (the production of glass and other mineral production and use).

Emissions from 2A2 Lime production are not estimated due to the lack of consistent activity data (lime production is located at four sites); those from 2A5 Asphalt roofing and 2A6 Road paving with asphalt are not estimated since no methodology is available.

Due to allocation problems, total emissions from mineral products (2A) are reported in category 2A7. Only particulate matter emissions could be reported separately in the category 2A1 Cement production. Particulate matter emissions in this category are derived from the environmental reports of the companies.

Other mineral products, including non-fuel mining and construction (2A7)

Category 2A7 is a key source for SO₂ and Pb emissions. The SO₂ emissions reported in this category originates in glass production and production of roof tiles. The Pb emissions in this category are from glass production only. The SO₂ and lead emissions from glass production are calculated with the extrapolation method (1), see section 5.1, based on m² glass produced by individual companies and total glass production, as reported by CBS. The SO₂ emission from the production of roof tiles is based on production statistics and a specific emission factor (method 3)

Category 2A7 is a key source for the following components (% of national total):

SO_x	(1.5%)	SBI 261 Glass production (1Gg) and SBI 264 production of roof tiles (0.9 Gg in 2004)
TSP	(2.4%)	
PM₁₀	(3.1%)	
PM_{2,5}	(1.9%)	
Pb	(8.6%)	SBI 261 Glass production
DIOX	(0.9%)	

4.2 Chemical industry (2B)

The PER comprises emissions related to three source categories as belonging to this category:

- 2B1 Ammonia production (SBI 24.15 ‘Manufacture of artificial fertilisers’)
- 2B2 Nitric acid production (included in SBI 24.1 ‘Manufacture of basic organic chemicals’)
- 2B5 Emissions from ‘Other chemical product manufacture’:
 - Manufacture of chemicals for agricultural use;
 - Manufacture of other chemical products (glue, photo chemicals, pharmaceuticals, fibres, paint and ink, soap and detergents).

Adapic acid (2B3) and calcium carbide (included in 2B4) are not produced in the Netherlands.

Emissions are not reported under 2B2 Nitric acid production (only the greenhouse gas N₂O is reported here). Due to allocation problems, all emissions from the chemical industry (2B) are reported in category 2B5.

Other chemical industry (2B5)

Category 2B5 is a key source for the following components (% of national total):

NMVO	(5.6%)
TSP	(4.3%)
PM₁₀	(2.2%)
PM_{2.5}	(2.1%)
Pb	(5.4%)
Hg	(22.3%)
Cd	(46.9%)

All emissions are calculated by extrapolation of activity data and emissions of the individual companies to totals in the subsectors, using production volume or production value as a proxy.

4.3 Metal production (2 C)

The national inventory of the Netherlands comprises emissions from Iron and steel production and aluminium production. The Netherlands has one integrated iron and steel plant (Corus, formerly known as Hoogovens). Integrated steelworks convert iron ores into steel by means of sintering, producing pig iron in blast furnaces and converting pig iron to steel in basic oxygen furnaces. For the purpose of the inventory, emissions from integrated steelworks are estimated for these three processes, as well as for some other minor processes. Emissions from sintering are included in 1A. A portion of the coke oven gas and blast/oxygen furnace gas produced during these processes is sold to a nearby power plant to be used as fuel. These emissions are included in category 1B. Aluminium is produced at two primary aluminium smelters (Pechiney and Aldel).

The above-mentioned companies report their emissions in environmental reports. Extrapolations to total emissions of the sector (method 1, see section 5.3) are very small, except for PM₁₀ and PM_{2.5}. CO, Pb, Cd, dioxins and PAH emissions in Category 2C are covered for more than 96% by individual registration. For PM₁₀, this is 83%.

Category 2C is a key source for the following components (% of national total):

CO	(9.5%)
TSP	(9.1%)
PM₁₀	(5.7%)
PM_{2.5}	(4.9%)
Pb	(59.7%)
Hg	(21.2%)
Cd	(43.1%)

4.4 Pulp and paper production (2D1)

Particulate matter emissions in this category are derived from the environmental reports of the companies and completed with calculations of specific emission factors multiplied by production figures (supplied by CBS).

Category 2D1 is a key source for the following component (% of national total):

PM₁₀	(1.0%)
------------------------	--------

4.5 Food and drink production (2D2)

NMVOC emissions in this category are derived from the environmental reports of the companies. Particulate matter emissions are calculated by multiplying specific emission factors by production figures (supplied by CBS).

Category 2D2 is a key source for the following components (% of national total):

NMVOC	(2.8%)
TSP	(3.8%)
PM₁₀	(3.4%)
PM_{2,5}	(0.9%)

4.6 Other production (2G)

See 2D2 Food and drink production for NMVOC. Most of the PAH emissions are emitted in the electrical engineering sector. The emission figures come from the environmental reports of the involved companies.

Category 2G is a key source for the following components (% of national total):

NMVOC	(8.2%)
TSP	(15.5%)
PM₁₀	(8.6%)
PM_{2,5}	(1.5%)
PAH	(13.6%)

5 Solvents and product use (3)

5.1 Paint application (3A)

VOC emissions from paint are calculated from national paint sales statistics (defined as paint that is both produced and sold in the Netherlands), provided by the Netherlands Association of Paint Producers VVVF (VVVF, 1999) and from paint imports, estimated by VVVF. The VVVF (through its members) directly monitors VOC in paint, while an assumption of the VVVF is used for the VOC in imported paint. Estimates have also been made for paint-related thinner use and the (reduction) effect of afterburners. For more information, see the protocol 'Calculation of VOC emissions from paint in the Netherlands' (Peek, 2007).

Category 3A is a key source for the following components (% of national total):

NMVOG	(14.8%)
-------	---------

5.2 Degreasing and dry cleaning (3B)

No key sources in this category.

5.3 Chemical products, manufacture and processing (3C)

No key sources in this category.

5.4 Other, including products containing HMs and POPs (3D)

The most relevant emission sources in this category are the companies for storage and transfer of oil products, chemicals and dry bulk commodities. These activities are major sources for NMVOG, TSP and PM₁₀ emissions in the Netherlands. The emissions are estimated on the basis of data from the sector (environmental reports from large companies).

This category also includes the emissions from the use of creosoted wood products (PAH) and dioxin emissions from PCP treated wood. The emission is estimated using a specific Dutch method

Based on Bremmer et al (1993) dioxin emissions by wooden house frames are determined for 1990. Since PCP has been banned from 1989, a linear reduction of dioxin emission has been assumed from ca 25 g I-TEQ in 1990 to ca 20 g I-TEQ in 2000.

Category 3D is a key source for the following components (% of national total):

NMVO	(13.9%)
TSP	(2.3%)
PM₁₀	(2.9%)
DIOX	(64.4%)
PAH	(44.8%)

6 Agriculture (4)

The sector Agriculture is a major source category for ammonia and particulate matter emissions. See section 5.2.3 on emission trends. For ammonia emission from agriculture two different sources are distinguished: animal manure and synthetic fertiliser. The main source of agricultural particulate matter emissions is formed by animal housing systems. A description of the calculation method is provided in Van der Hoek (2002).

Ammonia emissions from animal manure

Ammonia emissions from animal manure are calculated using the Manure and Ammonia model developed by the Dutch Agricultural Economics Research Institute (LEI). Input data for this model are divided into general and specific. General input data are animal numbers taken from the annual agricultural census. Specific input data concern the nitrogen and phosphate excretion by different animal categories and the ammonia volatilisation rates from animal housing systems and soil application systems for animal manure. The average nitrogen excretion per animal category is calculated annually as the difference between absorbed nitrogen from feeding and the capture in animal products. This so-called 'balance' - method takes into account annual changes in food consumption, food nitrogen content, etc. The excreted nitrogen partly volatilises as ammonia in stables, in pasture, during storage and during application to the soil. The share of housing and manure application systems with low ammonia volatilisation rates is taken into account. The rate of volatilisation of ammonia from animal manure depends on such aspects as the nitrogen content of the manure, the chemical balance between ammonia and ammonium in the manure and, finally, the contact surface manure – air and the exposure time.

Ammonia emissions from synthetic fertiliser

Ammonia emissions from synthetic fertiliser are calculated using data on the amounts of applied nitrogen fertiliser. These data are recorded by LEI and reported in Landbouwcijfers (Agricultural Data); these are also available via www.lei.wur.nl. Several types of nitrogen - each with their own specific ammonia emission factor fertiliser- are distinguished.

Particulate matter emissions from Agriculture

The main source for particulate matter emissions from agriculture are animal housing systems. Some other smaller sources include application of synthetic fertiliser, application of pesticides, supply of concentrates, hay making and harvesting of arable crops. The general input data for calculating the emissions from animal housing systems are animal numbers taken from the annual agricultural census. The share of poultry in free range housing systems with relative high emission factors is taken into account in these calculations. For several sources country-specific emission factors are available (Chardon and Van der Hoek, 2002).

6.1 Dairy cattle (4B1A)

Ammonia emission from dairy cattle (adult female cows) is calculated by multiplying the activity number with the emission factor. For dairy cattle the activity number is based on the animal number count from the annual agricultural census. Emission factors are calculated from the excretion and volatilisation rates for dairy cows.

Excretion rate

The excretion rate depends on feed intake and milk yield. Distinction is made between type of feed in two regions in the Netherlands and for two periods (summer and winter period).

Volatilisation rate

There are four different sources for ammonia emissions from animal manure: animal housing (manure production and storage), outside storage facilities (manure storage), meadow (manure production) and soils (manure application). For this reason also four different volatilisation rates are distinguished. Approximately 40% of dairy cattle ammonia emissions are derived from application of manure to the soil. Another 40% derives from animal houses and the rest is from animals grazing and from storage of manure.

Category 4B1a is a key source for the following component (% of national total):

NH ₃	(28.2%)
PM ₁₀	(2.2%)

6.2 Non-dairy cattle (4B1b)

Emission data for non-dairy cattle is based on the non-dairy cattle number count from the agricultural census and on emission factors calculated from excretion and volatilisation rates for these specific cows. There is, however, no distinction made between young and full-grown animals. Young dairy cattle are also included in this category.

Category 4B1b is a key source for the following component (% of national total):

NH ₃	(16.2%)
-----------------	---------

6.3 Swine (4B8)

Emission data is based on agricultural data from the agricultural census and on emission factors calculated from excretion and volatilisation rates for two specific types of pigs. Distinction is made between animals for meat production (fattening pigs) and animals used for breeding (breeding sows). Three types of volatilisation rates are distinguished for the two swine categories: the animal house, manure storage and soil application. Housing systems with high and low ammonia emission are also distinguished.

Emission of particulates comes mainly from skin, manure, feed and mulch particles ventilated from animal housing and have been calculated using an average emission factor per pig, deducted from the PM₅ (respirable fraction of PM) emission factor.

Category 4B8 is a key source for the following component (% of national total):

NH₃	(23.3%)
TSP	(4.9%)
PM₁₀	(6.2%)
PM_{2.5}	(2.1%)

6.4 Poultry (4B9)

Ammonia emission data is based on agricultural data from the agricultural census and on emission factors calculated from excretion and volatilisation rates for two specific types of poultry. Distinction is made between animals for meat production (broilers) and animals for egg production (laying hens). For both poultry categories, three types of volatilisation rates are distinguished: for the animal house, manure storage and soil application. Five different housing systems are distinguished for laying hens.

Emission of particulates comes mainly from skin, manure, feed and mulch particles ventilated from animal housing; these have been calculated using an average emission factor per chick, deducted from the PM₅ (respirable fraction of PM) emission factor. A distinction is made between free-range housing systems (with high particulate matter emission factors) and housing systems based on cages (with a low particulate matter emission factor).

Category 4B9 is a key source for the following component (% of national total):

NH₃	(12.5%)
TSP	(10.0%)
PM₁₀	(12.6%)
PM_{2.5}	(4.4%)

6.5 Other agricultural emissions (4G)

Emission data is based on amounts of different types of synthetic fertiliser and specific emission factors.

Category 4G is a key source for the following component (% of national total):

NH₃	(9.4%)
-----------------------	--------

7 Waste (6)

7.1 Waste incineration (6C)

The combustion emissions from waste incineration are included in the category 1A1a because nearly all the energy from waste incineration is converted into electricity. The emissions from crematoria are reported in this category. Mercury (Hg) emissions are estimated on the basis of the number of corpses cremated and a average amalgam content.

Category 6C is a key source for the following components (% of national total):

Hg	(9.4%)
DIOX	(0.9%)

7.2 Other waste (6D)

The emissions from the waste sector are reported in this category and coded as process emissions in the database. For historical reasons this category used to contain some of the emissions from waste incineration. This has been corrected in the latest submission, and emissions have been transferred to the 1A1a category.

No key sources in this category.

8 Other (7)

The emissions from burning candles, smoking of cigarettes and lighting of fireworks are reported in this category. This also includes the emissions of NH₃ from human perspiration. Please note that the Netherlands include this NH₃ sources in the national total, whereas other parties do not. There is no clear guidance yet whether or not these emissions should be included in the national total for NH₃.

Category 7 is a key source for the following components (% of national total):

NH₃	(3.8%)
TSP	(2.9%)
PM₁₀	(3.6%)
PM_{2,5}	(6.3%)

9 Recalculations and other changes

9.1 Recalculations compared to 2006 submission

In the preparation of the 2007 submission, some methodological improvements were included in the PER system. This led to some recalculations, aiming at the improvement of the accuracy of the emission data for the Netherlands. The emissions for 2004 were also updated on the basis of more accurate statistical information. Some emissions were reallocated. Compared to the 2006 submission, the following major changes can be distinguished:

- NMVOC emissions from Traffic and Transport (evaporation from passenger cars and small vans) were recalculated on the basis of lower emission factors derived from the COPERT model (emissions decreased over the whole time series by 20 – 25 Gg (about 25 Gg in 2004, compared to the reported emission from the 2006 submission).
- PM₁₀ emissions from brake wear were recalculated. Research showed that a substantive part of the emitted particles belonged to the coarse fraction (in the past, all emissions were calculated as PM₁₀, which led to a reduction of PM₁₀ emissions in the order of 0.6-0.8 Gg over the time series (0.7 Gg for 2004).
- Ammonia emissions from fertilisers have now been calculated, taking into account more detailed information of the different types of synthetic fertilisers. This led to an increase in emissions over the time series of 0.5 Gg in 1990 up to 4.3 Gg in 2003. This methodological change had already been applied to the year 2004 in the former submission. Comparing 2004 emission data between the 2006 submission and the 2007 submission, noted differences can be clarified by the availability of updated statistical data.
- Emissions from the oil and gas sector were recalculated on the basis of updated information received from the sector in their Annual Environmental Report. This resulted in a reduction of NO_x, SO₂ and NMVOC emissions over the whole time series. Compared to the 2006 submission, 2004 emissions for NMVOC decreased by about 7 Gg; SO₂ emissions decreased about 2 Gg and NO_x emissions increased about 2 Gg.
- The 2004 NO_x emissions decreased by about 6 Gg compared to the 2006 submission. The reason for this decrease is the availability of updated statistical data.
- Changing definitions led to different allocations: NMVOC emissions in category 1A1b (petroleum refining) decreased by some 2 Gg in 2004, while the emissions in category 1B2a iv (refining) increased by the same order of magnitude.
- All other (in most cases small) changes result from the availability of better (more detailed/accurate) statistical information.

9.2 Developments in emission insights and estimates

Since committing to the goals of the NEC ceiling directive in 2001, the insights in historical emissions and future estimates for 2010 have changed. Both the EU and the UNECE/CLRTAP state that countries should report their emissions according to best knowledge, even when this leads to other policy demands than accounted for at the time of drawing up the goals. The EU is clearly aware that evolving knowledge on emissions could lead to extra efforts, but still has not stated how this will be dealt with when assessing the NEC ceilings directive. Presenting the differences between the emissions calculated with old and new insights can facilitate the discussion on this. Evolving insights can be listed as follows:

- insights in scenario assumptions on future economic, social and technological developments;
- insights through evolving scientific knowledge, such as new insights into emission factors of sources or a changed method (model) for making an inventory of emissions;
- insights by formal adaptation within a sector definition through which a certain source is moved to another sector;
- insights through new policies.

The starting point for the Dutch analysis was the GC (i.e. Global Competition scenario) estimates from Environmental Outlook 5 for the year 2000 (MV5-GC), [RIVM, 2000]. These estimates were also used as a base for negotiations on the ceilings of the NEC directive. Table 9.1 gives the remaining policy efforts according to the insights in 2000.

Table 9.1. Estimates for 2010 according to the MV5 GC-scenario per sector (million kg) from 2000

Sector	SO ₂	NO _x	NH ₃	NMVOC	PM ₁₀
Industry, Energy and Refineries	55	76 ⁴⁾	4.1	84,7	10.9
Transport	15	180	NE ¹⁾	53	10.2
Trade, Services, Government, Construction	3 ²⁾	9 ⁵⁾	0.7	38.3	0.7
Consumers	2 ²⁾	15.4	7.3	37,8	6.5
Agriculture	2 ²⁾	9.8	137 ³⁾	2.4	1 ¹⁾
Total	72	290	149	216	28
NEC ceiling	50	260	128	185	-
Remaining policy effort in 2000	22	30	21	31	-

¹⁾ Not estimated

²⁾ Including consumers and agriculture

³⁾ EC scenario

⁴⁾ Including waste

⁵⁾ Excluding waste

Table 9.1 presents a detailed synopsis on the main evolving insights per sector per NEC substance introduced since the MV5-GC estimates from 2000. The most important new insights concern the traffic emission factors (cycle bypassing in road traffic), scenario assumptions in traffic (lower volume expectations for freight on road and water), exclusion of the international sea-ship emissions for NEC compliance, the lower volatilisation of NH₃ from manure and lower fugitive NMVOC emissions from gasoline-fuelled vehicles. A summary of these evolving insights is given in Table 9.2.

Table 9.2. Summary of new insights into emission estimates (million kg for 2010 according to type of evolving insight observed in the period 2000-2006)

Type of evolving insight	SO ₂	NO _x	NH ₃	NMVOC	PM ₁₀
1) Scenario assumptions	+4	-6	+4	-5 to -26	+14
2) Emission factors	-	+12	-11	-13	
3) Formal definitions	-9	-17	0	0	0
Other ¹⁾	-1	+4	+4	+2	+1
Total evolving insights	-6	-7	-3	-16 to -37	+15
New policy in 2000-2006	-	-6	-20	-37 to -16	-2
Total	-6	-13	-23	-53	+13

- 1) Not every evolving insight contributes to one of the four distinct types as good as another. Furthermore, through interdependencies between new insights the total within a sector can deviate somewhat from the sum of parts.

From Table 9.2 it appears that when all evolving insights between 2000 and 2006 (new policy excluded) are totaled over the sectors, the remaining policy efforts for all NEC substances have decreased since 2000. For SO₂ this totals -6 million kg, NO_x -7 million kg, NH₃ -3 million kg and NMVOC -16 to -37 million kg. The effects of new policy are presented separately in Table 9.2 since new policy concerns the sphere of influence from the Dutch government.

In February 2007, the Netherlands submitted a revised prognosis to UNECE (Table 9.3)

Table 9.3. Submission of the Netherlands

Pollutant:	UNIT	Current legislation projections			Current reduction plans		
		2010	2015	2020	2010	2015	2020
Sulphur oxides (SO _x as SO ₂)	Gg	51 ^(a)	55	57	49	51	53
Nitrogen oxides (NO _x as NO ₂)	Gg	262	233	219	261	226	203
Non-methane volatile organic compounds (NMVOC) ^(b)	Gg	162	NA	NA	NA	NA	NA
Ammonia (NH ₃)	Gg	125	135	145	123	133	143

(a) According to the Environmental Assessment Agency (MNP), the projected emission in 2010 for SO₂ is 53 Gg.

Compared to the prognosis submitted in 2006 (as presented above), differences can be clarified as follows:

- The 2007 prognosis includes a number of new emission insights that have not yet been included in the national PER.
- During 2006, new policies became effective.
- Some errors (double counting) have been corrected.

New emission insights:

- The new model VERSIT+ is applied to calculating emissions for the traffic and transport sector (Smit et al., 2006). NO_x emissions decrease by 17.6 Gg; PM₁₀ emissions decrease by 0.2 Gg and NH₃ emissions decrease by 0.2 Gg (all in 2010). New insights with respect to NMVOC have not yet been included in the prognosis;
- Emissions from companies for storage and transfer of oil products, chemicals and dry bulk commodities were recalculated on the basis of new insights: PM₁₀ emissions decrease by 0.1 Gg in 2010.

New policies:

- A new agreement between government and energy producers and refineries has lead to a projected SO₂ reduction in 2010 of 13 Gg; PM₁₀ emissions decrease by 0.5 Gg in 2010.

Error correction:

- Some double counting with respect to consumer emissions of PM₁₀ leads to a reduction in projected emissions fro 2010 of 1.4 Gg

9.3 Improvements

In the IIR 2006 - and in response to the review of the 2006 submission to the LRTAP Convention - the Netherlands reported a number of planned improvements for the following submissions. At the moment, the status of these improvements is given below.

Improvements as announced in the IIR 2006:

- *Improvement of consistency in the use of notation keys:* an effort was made to improve the consistency in the use of notation keys for the NFR submission of 2007.
- *Removal of inconsistencies in the source allocation over the years. For historical reasons, for example, the emissions from waste incineration could not be allocated in IA1a for all years. The category 3D currently holds the company emissions for storage and transfer of oil products, chemicals and dry bulk commodities. These sources could better be allocated in category 2 (industry):* inconsistencies in source allocation have been removed in the 2007 submission.
- *Time series consistency. In the current submission system, the Netherlands only submits data for 1990, 1995, 2000, 2002, 2003 and 2004. The years in between are not reported because they are not recalculated. In the next submission, these years will be re-submitted using an interpolation of the data based on the data (years) recalculated every year:* the 2007 submission contains a complete time series, where necessary, on the basis of interpolation.

Current improvements to the Informative Inventory Report

- *Quantitative effect of recalculation for abovementioned inventory years and for projected years:* see section 7.1 and 7.2.
- *Overview of new or expected evolving insights in the emission data:* see section below.

- *Documentation of the emission factors (e.g., English translation of the 'Methodological report for Traffic')*. Documents describing the methodologies applied in the PER are available on the website www.prtr.nl. At this time, improved protocols are developed, especially for SO₂, NO_x and NMVOC.
- *Documentation of methods for all key sources for PM₁₀ and PM_{2.5}*: this documentation is still under development.
- *Speciation profiles for NMVOC and particulate matter*: TNO has been working on this speciation on the basis of a literature search. The report is still in preparation.

Planned improvements to the inventory

A number of new insights/ improvements will be included in the upcoming inventory. As mentioned above, some of these new insights have already been included in the prognosis submitted by the Netherlands in February 2007:

(1) results of a new model for calculating the emissions from the traffic and transport sector will be included. NO_x, PM₁₀ and NMVOC emissions will decrease by about 5%. CO emissions will increase by about 5%. New activity data for this sector will also be included.

(2) new activity data for high efficiency boilers in households have become available. As a result, NO_x emissions will decrease.

(3) gas engines in horticulture and arable farming.

Faster market penetration of gas engines and increasing numbers and use of Biox power stations. These changes at least influence the projected emissions; historical effect remains to be explored.

(4) PAH emissions from the application of creosote oil.

Reduction of B(a)P emission by creosote oil application since 2000.

(5) new insights into the calculation of PM_{2.5} emissions have become available.

Based on literature research, the calculation of PM_{2.5} emissions will be further improved.

(6) individual PAH emissions

Based on research in 2007/2008, the Netherlands expects to be able to report the individual PAH emissions, instead of total PAH emissions.

(7) documentation

The documentation of the methodologies applied to the Netherlands PER system will be further improved and will be available on www.prtr.nl.

Furthermore, studies will be performed to further improve the methodologies applied to the LRTAP compounds; especially PM (PM₁₀ as well as PM_{2.5}), heavy metals and Persistent Organic Pollutants. The energy sector, households, waste sector and the commercial/institutional sector will be dealt with first.

References

- Alkemade G.E.M., C.J. Peek and P.G. Ruysenaars (2005). Prioritaire stoffen in de EmissieRegistratie. Emissies naar lucht, kwaliteit en aanbevelingen. MNP report 500055003/2005, Netherlands Environmental Assessment Agency, Bilthoven (in Dutch).
- Asselt, M.B.A. van (2000). Perspectives on uncertainty and risk-the PRIMA approach to decision-support, PhD thesis, Universiteit Maastricht.
- Bremmer, H.J., L.M. Troost, G. Kuipers, J. de Koning, A.A. Sein (1993), Emissies van dioxinen in Nederland, RIVM rapportnummer 770501003, National Institute for Public Health and the Environment RIVM, Bilthoven (in Dutch).
- Chardon, W.J. and K.W. van der Hoek (2002) Berekeningsmethode voor de emissie van fijn stof vanuit de landbouw, RIVM rapportnummer 773004014, National Institute for Public Health and the Environment RIVM, Bilthoven (in Dutch).
- DERA (1999). Second revised edition of the ICAO Aircraft Engine Exhaust Emissions Data Bank, DERA.
- EEA (2005). EMEP/CORINAIR Emission Inventory Guidebook – 2005. Technical report No 30. 2005/12/31. European Environment Agency. <http://reports.eea.eu.int/EMEPCORINAIR4/en>
- Eggleston, H.S. (1989). CORINAIR working group of emission factors for calculating 1985 emissions from road traffic Volume 1 ; methodology and emission factors. Consumer Protection and Nuclear Safety: Commission of the European Communities. Directorate-General Environment.
- EPA (1985) Environmental Protection Agency, Compilation of air pollution emission factors volume 2, Mobile sources, 4th edition.
- EMEP (2007) Inventory review 2007, Emission data reported to LRTAP Convention and NEC Directive, Technical report MSC-W 1/2007, EMEP.
- FAA (1996). Federal Aviation Administration, Federal Aviation Air Emissions Databank version 2.1., <http://www.faa.gov/>
- Gijlswijk, R. van, Coenen, P., Pulles, T. and Sluijs, J.P. van der, (2004), Uncertainty assessment of NO_x, SO₂ and NH₃ emissions in the Netherlands, TNO report R2004/100, TNO, Apeldoorn, Available: <http://www.rivm.nl/bibliotheek/digitaaldepot/TNOreportRa2004.pdf>.
- Harmelen, A.K. van, Denier van de Gon H.A.C., Kok H.J.G., Appelman W.J., Visschedijk A.J.H. and Hulskotte J.H.J. (2004). Particulate matter in the Dutch Pollutant Emission Register: State of Affairs. TNO report R2004/428. TNO Built Environment and Geosciences, Apeldoorn.
- Haskoning (2000), Diffuse emissies van fijn stof door (semi-)industriële activiteiten. HASKONING, Nijmegen (in Dutch).
- Hoek, K.W. van der, (2002), Uitgangspunten voor de mest- en ammoniakberekeningen 1999 tot en met 2001 zoals gebruikt in de Milieubalans 2001 en 2002, inclusief dataset landbouwemissies 1980 – 2001, RIVM rapportnummer 773004013, National Institute for Public Health and the Environment RIVM, Bilthoven (in Dutch).
- Hulskotte J.H.J. et al., (1999), Monitoringsystematiek openhaarden en houtkachels, TNO-MEP-R99/170, Apeldoorn (in Dutch).
- Hulskotte, J.H.J. and Koch, W.W.R. (2000), Emissiefactoren zeeschepen, TNO-MEP-R2000/221, Apeldoorn (in Dutch).
- Hulskotte, J.H.J., Bolt, E., Broekhuizen D. (2003), EMS-protocol emissies door binnenvaart: verbrandingsmotoren, RWS-AVV Ministry of Traffic, Public Works and Water Management, 's-Gravenhage (in Dutch).
- Hulskotte, J.H.J., Oonk, J., Roovaart, J.C. (2005), Emissieschattingen diffuse bronnen, Waterverontreiniging door motoremissies uit de recreatievaart, TNO-MEP, Apeldoorn (in Dutch).

- Klein, J.A.P., 1992. Luchtverontreiniging Emissies door Wegverkeer. Methodiek Vaststelling Emissiefactoren, Publikatiereeks Statistische Onderzoekingen M45, SDU/uitgeverij/CBS-publikaties, Den Haag (rapport opgesteld door het CBS in opdracht van Rijkswaterstaat).
- Klein, J.A.P., Hoen, A., Hulskotte, J.H.J., Duynhoven, N. van, Smit, R., Hensema, A. and D. Broekhuizen (2006). [Methods for calculating the emissions of transport in the Netherlands](#). Task Force Traffic and Transport of the National Emission Inventory.
- Koch W.W.R. (2001), Emission data for the Netherlands, 1998 and estimates for 1999. Report series monitoring of target groups no. 7., January 2001. Inspectorate for Environmental Protection, Department for Monitoring and Information Management. 's-Gravenhage.
- MNP (2005), Annual project plan for the Emission Inventory, Netherlands Environmental Assessment Agency. Bilthoven.
- MNP (2006), Annual project plan for the Emission Inventory, Netherlands Environmental Assessment Agency. Bilthoven.
- Most, P.F.J. van der, Loon, M.M.J. Van, Aulbers, J.A.W. and Daelen, H.J.A.M. van (1998), Methods for calculation of emissions to air and water (in Dutch). Report Series Emission Registration no. 44, July 1998, VROM-HIMH, The Hague.
- Ntziachristos, L., Z. Samaras (2000). COPERT III; Computer Programme to calculate emissions from road transport, methodology and emission factors (version 2.1), European Energy Agency (EEA), Copenhagen.
- Oonk et al. (2003), Emissiefactoren voor de binnenscheepvaart, TNO-rapport R2003/437 versie 2, Apeldoorn.
- Oorschot, M.M.P. van, Kraan, B.C.P., Brink, R.M.M. Van den, Janssen P.H.M. and Cooke, R.M. (2003), Uncertainty analysis for NO_x emissions from Dutch passenger cars in 1998. Applying structured expert elicitation and distinguishing different types of uncertainty, RIVM report 550002004, National Institute for Public Health and the Environment RIVM, Bilthoven.
<http://www.rivm.nl/bibliotheek/rapporten/550002004.html>.
- Peek, C.J. (2007) Protocol NMVOS emissies uit verfgbruik in Nederland, MNP Netherlands Environmental Assessment Agency, Bilthoven (in Dutch), In prep.
- Pulles, T., and Aardenne, J. van (2001), Good Practice Guidance for CLRTAP Emission Inventories, Draft chapter for the UNECE CORINAIR Guidebook on Emission Inventories, Available: <http://reports.eea.eu.int/EMEPCORINAIR4/en/BGPG.pdf> , 24 June, 2004]
- RIVM (1999), Measuring, Modelling and Uncertainty [in Dutch]. RIVM report 408129005 (main report and addendum), National Institute for Public Health and the Environment RIVM, Bilthoven.
- RIVM (2001), Environmental Balance 2000, RIVM report 251701051, National Institute for Public Health and the Environment RIVM, Bilthoven (in Dutch).
- Shareef, G.S., W.A. Butler, L.A. Bravo, M.B. Stockton (1988), Air emissions species manual, Vol. I: Volatile organic compound (VOC) species profiles. Radian Corp., EPA report 450/2-88-003a.
- Sluijs, J.P. van der, Risbey, J.S., and Ravetz, J. (2002), Uncertainty assessment of VOC emissions from paint in the Netherlands. Department of Science Technology and Society, Utrecht University, Available: <http://www.nusap.net>.
- Sluijs, J.P. van der, Klopogge, P., Risbey, J. and Ravetz, J. (2003), Towards a synthesis of qualitative and quantitative uncertainty assessment: applications of the numeral, unit, spread, assessment, pedigree (NUSAP) system, Paper for the International Workshop on Uncertainty, Sensitivity, and Parameter Estimation for Multimedia Environmental Modeling, August 19-21, 2003, Rockville, Maryland, USA, Available: <http://www.nusap.net>.
- Sluijs, J.P. van der, Craye, M., Funtowicz, S., Klopogge, P., Ravetz, J., and Risbey, J. (2005) Combining Quantitative and Qualitative Measures of Uncertainty in Model based Environmental Assessment: the NUSAP System, Risk Analysis, 25 (2). p. 481-492
- Soest-Vercammen, E.L.J. van, J.H.J. Hulskotte, D.C. Heslinga (2002): Monitoringsprotocol Bijschatting Stationaire NO_x-bronnen kleiner dan 20 MWth, TNO report R2002/042, TNO Milieu, Energie en Procesinnovatie, Apeldoorn (in Dutch).

Spakman, J. et al. (2003). [Method for calculating greenhouse gas emissions](#). Emission Registration Series/ Environmental Monitor No. 37b, March 2003: electronic update of no. 37, July 1997. VROM Inspectorate, The Hague.

Tak, C. van der (2000), Actualisatie T₀-emissies, report number 16196.620/2, MARIN, Wageningen

UNECE (2003), Guidelines for Estimating and Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution. Air Pollution Studies No. 15.

VROM (1993). Publikatiereeks Emissieregistratie. Emissiefactoren Vluchtige organische stoffen uit verbrandingsmotoren. no. 10, 's-Gravenhage (in Dutch).

VVVF (2004). Jaarrapportage Verf en drukinktgebruik in Nederland. Vereniging van verf- en drukinktfabrikanten, 's-Gravenhage (in Dutch).